

AD-A105 744

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS

F/8 12/1

MINIMAX ESTIMATION OF A MULTIVARIATE NORMAL MEAN UNDER A CONVEX-ETC(U)

AUG 81 P LIN: A MOUSA

N00014-80-C-0093

UNCLASSIFIED

FSU-STATISTICS-M591

NL

1 of 1  
482  
000711

SEARCHED	INDEXED
SERIALIZED	FILED
AUG 81 11 11 AM	
FBI - TAMPA	



END  
DATE  
FILMED  
11-81  
DTIC

ne Florida State University

Minimax Estimation of a Multivariate Normal Mean  
under a Convex Loss Function.

by  
Pi-Erh/Lin<sup>1</sup> and Amany/Mousa

FSU Statistics Report No. - M591  
ONR Technical Report No. - 153

11  
August, 1981

The Florida State University  
Department of Statistics  
Tallahassee, Florida 32306

15 FSU - STATISTICS - 11-16  
17A-153-ONR

15  
This work was supported by the Army, Navy and Air Force under Office of  
Naval Research Contract No. N00014-80-C-0093. Reproduction in whole or  
in part is permitted for any purpose of the United States Government.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Availability
A	Special

400 111

# Minimax Estimation of a Multivariate Normal Mean under a Convex Loss Function

by

Pi-Erh Lin<sup>1</sup> and Amany Mousa  
Florida State University

## Summary

Let  $\underline{X} = (X_1, \dots, X_p)' \sim N_p(\underline{\mu}, \Sigma)$  where  $\underline{\mu} = (\mu_1, \dots, \mu_p)'$  and  $\Sigma = \text{diag}(\sigma_1^2, \dots, \sigma_p^2)$  are both unknown and  $p \geq 3$ . Let  $(n_i - 2)w_i/\sigma_i^2 \sim \chi_{n_i}^2$ , independent of  $w_j$  ( $i \neq j = 1, \dots, p$ ). Assume that  $(w_1, \dots, w_p)$  and  $\underline{X}$  are independent. Define  $W = \text{diag}(w_1, \dots, w_p)$  and  $\|\underline{X}\|_W^2 = \underline{X}'W^{-1}Q^{-1}W^{-1}\underline{X}$  where  $Q = \text{diag}(q_1, \dots, q_p), q_i > 0, i = 1, \dots, p$ . In this paper, the minimax estimator of Berger and Bock (Ann. Statist. 4 (1976), 642-648), given by  $\hat{\delta}(\underline{X}, W) = [I_p - r(\underline{X}, W)\|\underline{X}\|_W^{-2}Q^{-1}W^{-1}]\underline{X}$ , is shown to be minimax relative to the convex loss  $(\hat{\delta} - \underline{\mu})'[\alpha Q + (1-\alpha)\Sigma^{-1}](\hat{\delta} - \underline{\mu})/C$ , where  $C = \alpha \text{tr}(\Sigma Q) + (1-\alpha)p$  and  $0 \leq \alpha \leq 1$ , under certain conditions on  $r(\underline{X}, W)$ . This generalizes the above-mentioned result of Berger and Bock.

---

<sup>1</sup>This work was supported by the Army, Navy and Air Force under Office of Naval Research Contract No. N00014-80-C-0093. Reproduction in whole or in part is permitted for any purpose of the United States Government.

AMS 1970 Subject Classifications. Primary 62C99, Secondary 62F10, 62H99.

Key words and Phrases. Unknown variances, convex combination of loss functions, risk function.

1. Introduction. Let  $\underline{X} = (X_1, \dots, X_p)'$  be a  $p$ -variate ( $p \geq 3$ ) random vector normally distributed with mean  $\underline{\mu} = (\mu_1, \dots, \mu_p)'$  and covariance matrix  $\Sigma = \text{diag}(\sigma_1^2, \dots, \sigma_p^2)$  where  $\sigma_i^2$ ,  $i = 1, \dots, p$ , are unknown. This paper obtains a class of minimax estimators for  $\underline{\mu}$  when the loss incurred in estimating  $\underline{\mu}$  by  $\underline{\delta}$  is given by

$$(1.1) \quad L(\underline{\delta}; \underline{\mu}, \Sigma) = (\underline{\delta} - \underline{\mu})' [\alpha Q + (1-\alpha)\Sigma^{-1}] (\underline{\delta} - \underline{\mu}) / C,$$

where  $0 \leq \alpha \leq 1$ ,  $C = \alpha \text{tr}(Q\Sigma) + (1-\alpha)p$ , and  $Q$  is a known  $p \times p$  diagonal matrix with diagonal elements  $q_i > 0$ ,  $i = 1, \dots, p$ . The loss (1.1) is a convex combination of two commonly used loss functions, namely

$$(1.2) \quad L_1(\underline{\delta}; \underline{\mu}, \Sigma) = (\underline{\delta} - \underline{\mu})' Q (\underline{\delta} - \underline{\mu}) / \text{tr}(Q\Sigma)$$

and

$$(1.3) \quad L_2(\underline{\delta}; \underline{\mu}, \Sigma) = (\underline{\delta} - \underline{\mu})' \Sigma^{-1} (\underline{\delta} - \underline{\mu}) / p.$$

The loss function  $L_1(\underline{\delta}; \underline{\mu}, \Sigma)$  may be used when the relative importance of the parameters to be estimated is reflected by a known set of weights represented by  $Q$ , while the loss  $L_2(\underline{\delta}; \underline{\mu}, \Sigma)$  represents the case when the relative importance is reflected naturally by the inverse of the covariance matrix of the variables. In the literature, statisticians tend to use either  $L_1(\underline{\delta}; \underline{\mu}, \Sigma)$  or  $L_2(\underline{\delta}; \underline{\mu}, \Sigma)$  as a loss function. In the convex loss (1.1) we have combined both viewpoints in such a way that the more  $\alpha$  is near 1 the more loss is to be assessed by  $L_1(\underline{\delta}; \underline{\mu}, \Sigma)$ , and vice versa.

It is clear that the maximum likelihood estimator,  $\underline{X}$ , is minimax with risk  $R(\underline{X}, \underline{\mu}) = E_{\underline{X}} L(\underline{X}; \underline{\mu}, \Sigma) = 1$ . Thus an estimator  $\underline{\delta}$  will be minimax under

loss (1.1) if and only if  $R(\underline{x}, \underline{\mu}) - R(\underline{\delta}, \underline{\mu}) \geq 0$  for all  $\underline{\mu}$  and  $\Sigma$ . In evaluating the difference in risk it is necessary to impose certain conditions on  $\Sigma$ . Specifically, the values or bounds of the trace of  $\Sigma^{-1}Q^{-1}$ , denoted by  $\text{tr}(\Sigma^{-1}Q^{-1})$ , and the minimum characteristic root of  $Q\Sigma$ , denoted by  $\text{ch}_{\min}(Q\Sigma)$ , will be assumed. Similar conditions have been noted by various authors. For example, Gleser (1976) shows that if a lower bound for  $\text{ch}_{\min}(Q\Sigma)$  is known then a family of minimax estimators for  $\underline{\mu}$  can be obtained under the loss function  $L_1(\underline{\delta}; \underline{\mu}, \Sigma)$ ; otherwise no estimator of the form  $[I_p - h(\underline{x}'W^{-1}\underline{x})Q^{-1}W^{-1}]\underline{x}$  can be minimax for  $\underline{\mu}$  unless  $h(u) = 0$  for almost all  $u \geq 0$  where  $W \sim W(n, \Sigma)$  independent of  $\underline{x}$  and  $h: R \rightarrow R$  is a real function satisfying certain conditions. In our case where both  $Q$  and  $\Sigma$  are diagonal matrices a sufficient set of conditions on  $\Sigma$  would be

$$(1.4) \quad \max_{1 \leq i \leq p} \{\sigma_i^2\} \leq 1/c_1 \quad \text{and} \quad \min_{1 \leq i \leq p} \{\sigma_i^2\} \geq 1/c_2$$

for some positive constants  $c_1$  and  $c_2$  ( $0 < c_1 \leq c_2 < \infty$ ). This set of conditions on  $\Sigma$  is not unreasonable in application.

2. Useful Lemmas. In establishing the minimaxity of an estimator of the mean of a multivariate normal distribution the following lemmas are very useful. They are presented here for ease of reference and without proof.

Lemma 2.1. [Stein (1974)]. Let  $Y \sim N(0, 1)$  and let  $g$  be an absolutely continuous function,  $g: R \rightarrow R$ . Then

$$E_Y[g'(Y)] = E_Y[Yg(Y)]$$

provided that all expectations exist and are finite.

Lemma 2.2. [Efron and Morris (1976)]. Let  $U \sim \chi_n^2$  and let  $g$  be as defined in Lemma 2.1. Then

$$E_U[Ug(U)] = nE_U[g(U)] + 2E_U[Ug'(U)]$$

provided that all expectations exist and are finite.

Corollary 2.2.1. Let  $U$  and  $g$  be as defined in Lemma 2.2. Let  $Z = cU/(n-2)$ ,  $c > 0$ , and  $h(Z) = g[(n-2)Z/c]$ . Then

$$E_Z[(n-2)Zh(Z)/c] = nE_Z[h(Z)] + 2E_Z[Zh'(Z)]$$

provided that all expectations exist and are finite.

Lemma 2.3. [Lehmann (1966)]. Let  $S$  be any random variable, and let  $p_1(S)$  and  $p_2(S)$  map the real line into itself. If  $p_1(S)$  and  $p_2(S)$  are either both nonincreasing in  $S$  or both nondecreasing in  $S$ , then

$$E_S[p_1(S)p_2(S)] \geq E_S[p_1(S)]E_S[p_2(S)].$$

Note that Lemmas 2.1 and 2.2 may be proved by integration by parts and the corollary by the indicated change of variable. Lemma 2.3 follows from a new concept of dependency, namely the positive quadrant dependence, introduced by Lehmann (1966). The above results will be employed in the proof of the main theorem in the next section.



3. Main Result. Let  $\underline{X}$  be a  $p$ -variate ( $p \geq 3$ ) random vector normally distributed with unknown mean  $\underline{\mu}$  and unknown covariance matrix  $\Sigma = \text{diag}(\sigma_1^2, \dots, \sigma_p^2)$ . Assume that  $(n_i - 2)w_i/\sigma_i^2 \sim \chi_{n_i}^2$ , ( $n_i > 2$ ),  $i = 1, \dots, p$ , where  $w_i$  and  $w_j$  ( $i \neq j = 1, \dots, p$ ) are mutually independent and are independent of  $\underline{X}$ . As in Berger and Bock (1976), define  $W = \text{diag}(w_1, \dots, w_p)$ ,  $T = \min_{1 \leq i \leq p} (\chi_{n_i}^2/n_i)$ , and  $\tau = \tau(n_1, \dots, n_p) = E(T^{-1})$ . In the following theorem we will obtain a class of minimax estimators for  $\underline{\mu}$  relative to the loss function given by (1.1).

Theorem. The estimator

$$(3.1) \quad \hat{\delta}(\underline{X}, W) = [I_p - r(\underline{X}, W) \|\underline{X}\|_W^{-2} Q^{-1} W^{-1}] \underline{X}$$

is minimax for  $\underline{\mu}$  relative to the loss function (1.1) with  $Q = \text{diag}(q_1, \dots, q_p)$ , where  $\|\underline{X}\|_W^2 = \underline{X}' W^{-1} Q^{-1} W^{-1} \underline{X}$ , provided that the following conditions are satisfied

$$(i) \quad 0 \leq r(\underline{X}, W) \leq \frac{2\{\alpha(p-2\tau) + (1-\alpha)[\text{tr}(\Sigma^{-1}Q^{-1}) - 2\tau \text{ch}_{\max}(\Sigma^{-1}Q^{-1})]\}}{\alpha + (1-\alpha)\text{ch}_{\max}(\Sigma^{-1}Q^{-1})} \text{ with}$$

$$\alpha(p-2\tau) + (1-\alpha)[\text{tr}(\Sigma^{-1}Q^{-1}) - 2\tau \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] \geq 0 \text{ and } 0 \leq \alpha \leq 1,$$

$$(ii) \quad r(\underline{X}, W) \text{ is nondecreasing in } |\underline{X}_i|, \quad i = 1, \dots, p,$$

$$(iii) \quad r(\underline{X}, W) \text{ is nonincreasing in } w_i, \quad i = 1, \dots, p,$$

$$(iv) \quad r(\underline{X}, W) \|\underline{X}\|_W^{-2} \text{ is nondecreasing in } w_i, \quad i = 1, \dots, p.$$

Proof. Write  $\hat{\delta} = \hat{\delta}(\underline{X}, W)$  and  $r = r(\underline{X}, W)$ . Let  $\Delta = R(\underline{X}, \underline{\mu}) - R(\hat{\delta}, \underline{\mu})$ .

Then

$$(3.2) \quad \begin{aligned} C\Delta &= E_{W, \underline{X}} \{ 2r \|\underline{X}\|_W^{-2} (\underline{X} - \underline{\mu})' [cQ + (1-\alpha)\Sigma^{-1}] Q^{-1} W^{-1} \underline{X} \\ &\quad - r^2 \|\underline{X}\|_W^{-4} \underline{X}' W^{-1} Q^{-1} [cQ + (1-\alpha)\Sigma^{-1}] Q^{-1} W^{-1} \underline{X} \} \\ &= 2[\alpha\Delta_1 + (1-\alpha)\Delta_2], \text{ say,} \end{aligned}$$

where

$$\Delta_1 = E_{W, X} [r \|X\|_W^{-2} (X - \mu)^T W^{-1} X - r^2 \|X\|_W^{-2} / 2]$$

and

$$\Delta_2 = E_{W, X} [r \|X\|_W^{-2} (X - \mu)^T \Sigma^{-1} Q^{-1} W^{-1} X - r^2 \|X\|_W^{-4} X^T W^{-1} Q^{-1} \Sigma^{-1} Q^{-1} W^{-1} X].$$

In the following we will find appropriate lower bounds for  $\Delta_1$  and  $\Delta_2$  which, in conjunction with Condition (i), will establish that  $\Delta \geq 0$ . But a lower bound for  $\Delta_1$  has been obtained by Berger and Bock (1976, Eq. (2.10)), namely

$$(3.3) \quad \Delta_1 \geq E_X \{ E_W (r \|X\|_W^{-2}) [p - 2\tau - E_W(r/2)] \}.$$

We will proceed to find a lower bound for  $\Delta_2$ . Let

$$(3.4) \quad \Delta_2 = \Delta_{21} + \Delta_{22}$$

where

$$(3.5) \quad \Delta_{21} = E_{W, X} [r \|X\|_W^{-2} (X - \mu)^T \Sigma^{-1} Q^{-1} W^{-1} X]$$

and

$$(3.6) \quad \begin{aligned} \Delta_{22} &= E_{W, X} [-(1/2) r^2 \|X\|_W^{-4} X^T W^{-1} Q^{-1} \Sigma^{-1} Q^{-1} W^{-1} X] \\ &\geq E_{W, X} [-(1/2) r^2 \|X\|_W^{-2} \text{ch}_{\max}(\Sigma^{-1} Q^{-1})]. \end{aligned}$$

Then, taking expectation with respect to  $X$  first and then  $W$ , and by an application of Lemma 2.1, we have

$$\begin{aligned}
(3.7) \quad \Delta_{21} &= E_{W, \underline{X}} \left[ \sum_{i=1}^p \frac{1}{\sigma_i q_i w_i} (r \|\underline{X}\|_W^{-2} X_i) \left( \frac{X_i - \mu_i}{\sigma_i} \right) \right] \\
&= E_{W, \underline{X}} \left[ \sum_{i=1}^p \frac{1}{q_i w_i} \frac{\partial}{\partial X_i} (r \|\underline{X}\|_W^{-2} X_i) \right] \\
&= E_{W, \underline{X}} \left[ \sum_{i=1}^p \left( \frac{r}{\|\underline{X}\|_W^2} - \frac{2r X_i^2}{\|\underline{X}\|_W^4 q_i w_i^2} + \frac{X_i}{\|\underline{X}\|_W^2} \frac{\partial r}{\partial X_i} \right) \right] \\
&\geq E_{W, \underline{X}} \left[ \frac{r}{\|\underline{X}\|_W^2} \sum_{i=1}^p \frac{1}{q_i w_i} - \frac{2r}{\|\underline{X}\|_W^4} \sum_{i=1}^p \frac{X_i^2}{q_i w_i^3} \right].
\end{aligned}$$

The last inequality follows from Condition (ii) since  $X_i (\partial r / \partial X_i) \geq 0$  for all  $i = 1, \dots, p$ . The first term in the right-hand side of (3.7) may be further evaluated by taking the expectation first with respect to  $W$  and then with respect to  $\underline{X}$ . Recall that  $(n_i - 2)w_i / \sigma_i^2 \sim \chi_{n_i}^2$ . Then, it follows from Corollary 2.2.1 with  $h(w_i) = r / (\|\underline{X}\|_W^2 w_i)$  that, for each  $i = 1, \dots, p$ ,

$$n_i E_{W_i} \left( \frac{r}{\|\underline{X}\|_W^2 w_i} \right) = \frac{n_i - 2}{\sigma_i^2} E_{W_i} \left( \frac{r}{\|\underline{X}\|_W^2} \right) - 2 E_{W_i} \left[ \frac{1}{\|\underline{X}\|_W^2} \frac{\partial r}{\partial w_i} + \frac{2r X_i^2}{\|\underline{X}\|_W^4 q_i w_i^3} - \frac{r}{\|\underline{X}\|_W^2 w_i} \right].$$

Therefore,

$$\begin{aligned}
(3.8) \quad E_{W, \underline{X}} \left( \frac{r}{\|\underline{X}\|_W^2 w_i} \right) &= E_{W, \underline{X}} \left[ \frac{r}{\|\underline{X}\|_W^2 \sigma_i^2} - \frac{4r X_i^2}{(n_i - 2) \|\underline{X}\|_W^4 q_i w_i^3} - \frac{2}{(n_i - 2) \|\underline{X}\|_W^2} \frac{\partial r}{\partial w_i} \right] \\
&\geq E_{W, \underline{X}} \left[ \frac{r}{\|\underline{X}\|_W^2 \sigma_i^2} - \frac{4r X_i^2}{(n_i - 2) \|\underline{X}\|_W^4 q_i w_i^3} \right].
\end{aligned}$$

The last inequality follows by Condition (iii). Now, substituting (3.8) into the last expression of (3.7), we have

$$\begin{aligned}
 (3.9) \quad \Delta_{21} &\geq E_{W, X} \left[ \frac{r}{\|X\|_W^2} \sum_{i=1}^p \frac{1}{\sigma_i^2 q_i} - \frac{4r}{\|X\|_W^4} \sum_{i=1}^p \frac{x_i^2}{(n_i-2)q_i w_i^3} - \frac{2r}{\|X\|_W^4} \sum_{i=1}^p \frac{x_i^2}{q_i w_i^3} \right] \\
 &= E_{W, X} \left[ \frac{r}{\|X\|_W^2} \sum_{i=1}^p \frac{1}{\sigma_i^2 q_i} - \frac{2r}{\|X\|_W^4} \sum_{i=1}^p \left( \frac{n_i}{n_i-2} \right) \frac{x_i^2}{q_i w_i^3} \right] \\
 &\geq E_{W, X} \left[ \frac{r}{\|X\|_W^2} \sum_{i=1}^p \frac{1}{\sigma_i^2 q_i} - \frac{2r}{\|X\|_W^2} \max_{1 \leq i \leq p} \left( \frac{n_i}{n_i-2} \frac{\sigma_i^2}{w_i} \frac{1}{\sigma_i^2 q_i} \right) \right] \\
 &\geq E_{W, X} \{ (r\|X\|_W^{-2}) [\text{tr}(\Sigma^{-1}Q^{-1}) - 2T^{-1} \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] \}.
 \end{aligned}$$

Substituting (3.9) and (3.6) into (3.4), we obtain

$$\begin{aligned}
 (3.10) \quad \Delta_2 &\geq E_{W, X} \{ (r\|X\|_W^{-2}) [\text{tr}(\Sigma^{-1}Q^{-1}) - 2T^{-1} \text{ch}_{\max}(\Sigma^{-1}Q^{-1}) - (r/2) \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] \} \\
 &\geq E_X \{ E_W(r\|X\|_W^{-2}) [\text{tr}(\Sigma^{-1}Q^{-1}) - 2\tau \text{ch}_{\max}(\Sigma^{-1}Q^{-1}) - E_W(r/2) \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] \}.
 \end{aligned}$$

The last inequality in (3.10) is obtained by an application of Lemma 2.3 since  $r\|X\|_W^{-2}$  is nondecreasing in  $w_i$  and  $[\text{tr}(\Sigma^{-1}Q^{-1}) - 2T^{-1} \text{ch}_{\max}(\Sigma^{-1}Q^{-1}) - (r/2) \text{ch}_{\max}(\Sigma^{-1}Q^{-1})]$  is also nondecreasing in  $w_i$  for  $i = 1, \dots, p$ , and since  $w_i$  and  $w_j$  ( $i \neq j$ ) are mutually independent. Hence, upon the substitution of (3.10) and (3.3) into (3.2), we conclude that

$$\begin{aligned}
 (3.11) \quad C\Delta/2 &= \alpha\Delta_1 + (1-\alpha)\Delta_2 \\
 &\geq E_X \{ E_W(r\|X\|_W^{-2}) \\
 &\quad \times \{ \alpha(p-2\tau) + (1-\alpha)[\text{tr}(\Sigma^{-1}Q^{-1}) - 2\tau \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] \\
 &\quad - [\alpha + (1-\alpha) \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] E_W(r/2) \} \}.
 \end{aligned}$$

By Condition (i), it is clear that the last expression of (3.11) is non-negative for all  $\underline{\mu}$  and  $\Sigma$  and hence  $\hat{\delta}$  is minimax proving the theorem.

Remarks. (1) Unless Condition (i) is satisfied, the theorem is vacuous. Thus it is required not only that  $\text{tr}(\Sigma^{-1}Q^{-1})$  and  $\text{ch}_{\max}(\Sigma^{-1}Q^{-1})$  be known but that, for  $0 \leq \alpha \leq 1$ ,

$$(3.12) \quad \alpha(p-2\tau) + (1-\alpha)[\text{tr}(\Sigma^{-1}Q^{-1}) - 2\tau \text{ch}_{\max}(\Sigma^{-1}Q^{-1})] \geq 0.$$

On the other hand, if there exists positive constants  $c_1$  and  $c_2$  ( $0 < c_1 \leq c_2 < \infty$ ) such that (1.4) holds or, equivalently,

$$\text{ch}_{\min}(\Sigma^{-1}) \geq c_1, \quad \text{ch}_{\max}(\Sigma^{-1}) \leq c_2$$

and

$$p \geq 2\tau[\alpha + (1-\alpha)c_2 \text{ch}_{\max}(Q^{-1})]/[\alpha + (1-\alpha)c_1 \text{ch}_{\min}(Q^{-1})],$$

then the theorem remains true with Condition (i) replaced by

$$(i') \quad 0 \leq r(\underline{X}, W) \leq \frac{2\{\alpha(p-2\tau) + (1-\alpha)[c_1 \text{tr}(Q^{-1}) - 2\tau c_2 \text{ch}_{\max}(Q^{-1})]\}}{\alpha + (1-\alpha)c_2 \text{ch}_{\max}(Q^{-1})}$$

(2) Berger and Bock (1976) show that the estimator  $\hat{\delta}(\underline{X}, W)$  given by (3.1) is minimax for  $\underline{\mu}$  under the loss  $L_1(\hat{\delta}; \underline{\mu}, \Sigma)$  and that there exists a large positive integer  $N$  such that, for all  $n_i \geq N$ ,  $p-2\tau > 0$  and  $p \geq 3$ . Furthermore, they provide a formula for calculating  $\tau$  when the  $n_i$ 's are even. In this paper, we employ their method of proof to establish our result. It should be noted that our result reduces to theirs when  $\alpha = 1$ .

### References

- Berger, J.O. and Bock, M.E. (1976). Combining independent normal mean estimation problems with unknown variances. Ann. Statist. 4, 642-648.
- Efron, B. and Morris, C. (1976). Families of minimax estimators of the mean of a multivariate normal distribution. Ann. Statist. 4, 11-21.
- Gleser, L.J. (1976). Minimax estimation of a multivariate normal mean with unknown covariance matrix. Technical Report No. 460, Purdue University.
- Lehmann, E.L. (1966). Some concepts of dependence. Ann. Math. Statist. 37, 1137-1153.
- Stein, C. (1974). Estimation of the parameters of a multivariate normal distribution, Part I, Technical Report No. 63, Stanford University.

## SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU No. M591 ONR No. 153	2. GOVT ACCESSION NO. <i>AD-A205744</i>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and subtitle) Minimax Estimation of a Multivariate Normal Mean under a Convex Loss Function	5. TYPE OF REPORT & PERIOD COVERED Technical Report	6. PERFORMING ORG. REPORT NUMBER FSU Statistics Report M591
7. AUTHOR(s) Pi-Erh Lin and Amany Mousa	8. CONTRACT OR GRANT NUMBER(s) ONR No. N00014-80-C-0093	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Statistics and Probability Program Arlington, Virginia 22217	12. REPORT DATE August, 1981	13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE

## 16. DISTRIBUTION STATEMENT (of this report)

Approved for public release: distribution unlimited

## 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)

## 18. SUPPLEMENTARY NOTES

## 19. KEY WORDS

Unknown variances, convex combination of loss functions, risk function.

## 20. ABSTRACT (Continue on reverse side if necessary and identify by block number)

Let  $\underline{X} = (X_1, \dots, X_p)' \sim N_p(\underline{\mu}, \Sigma)$  where  $\underline{\mu} = (\mu_1, \dots, \mu_p)'$  and  $\Sigma = \text{diag}(\sigma_1^2, \dots, \sigma_p^2)$  are both unknown and  $p \geq 3$ . Let  $(n_i - 2)w_i/\sigma_i^2 \sim \chi_{n_i}^2$ , independent of  $w_j$  ( $i \neq j = 1, \dots, p$ ).

Assume that  $(w_1, \dots, w_p)$  and  $\underline{X}$  are independent. Define  $W = \text{diag}(w_1, \dots, w_p)$  and  $\|\underline{X}\|_W^2 = \underline{X}' W^{-1} Q^{-1} W^{-1} \underline{X}$  where  $Q = \text{diag}(q_1, \dots, q_p)$ ,  $q_i > 0$ ,  $i = 1, \dots, p$ . In this paper, the minimax estimator of Berger and Bock (Ann. Statist. 4 (1976), 642-648), given by  $\hat{\underline{X}}(\underline{X}, W) = [I_p - r(\underline{X}, W) \|\underline{X}\|_W^{-2} Q^{-1} W^{-1}] \underline{X}$ , is shown to be minimax relative to the convex loss  $(\hat{\underline{X}} - \underline{\mu})' [\alpha Q + (1-\alpha)\Sigma^{-1}] (\hat{\underline{X}} - \underline{\mu}) / C$ , where  $C = \alpha \text{tr}(\Sigma Q) + (1-\alpha)p$  and  $0 \leq \alpha \leq 1$ , under certain conditions on  $r(\underline{X}, W)$ . This generalizes the above-mentioned result of Berger and Bock.